

CLAIMS

What is claimed is:

1. A method of identifying a compound capable of modulating the activity of Cathepsin
5 Z in a cell, comprising the steps of:
 - (a) measuring said cell's base level of Cathepsin Z activity in the absence of a candidate compound;
 - (b) introducing said candidate compound; and
 - (c) measuring said cell's level of Cathepsin Z activity in the presence of said
10 candidate compound.
2. The method of Claim 1 wherein said cell's level of Cathepsin Z activity is measured by measuring antigen presentation.
- 15 3. The method of Claim 2, wherein said cell's level of antigen presentation is measured by measuring autologous T-cell response to tetanus toxin.
4. The method of Claim 2, wherein said cell's level of antigen presentation is measured by measuring said cell's capacity to present quenched FITC-ovalbumin.
- 20 5. The method of Claim 1 wherein said cell is an antigen-presenting cell.
6. The method of Claim 5 wherein said antigen-presenting cell is selected from the group consisting of dendritic cell precursor, immature dendritic cell, and mature dendritic
25 cell.
7. A compound capable of modulating the activity of Cathepsin Z identified according to the method of Claim 1.
- 30 8. A pharmaceutical for treating an inflammatory disease comprising the compound of Claim 7 and a pharmaceutically acceptable excipient.

9. A method for treating an autoimmune disease comprising the step of administering the pharmaceutical of Claim 8.

10. The method of Claim 9 wherein said autoimmune disease is selected from the group consisting of rheumatoid arthritis and multiple sclerosis.